

Higher Derivatives

11.3



Introduction

The derivative, $\frac{dy}{dx}$, is more expressly called the **first derivative** of y . By differentiating the first derivative, we obtain the **second derivative**; by differentiating the second derivative we obtain the **third derivative** and so on. These second and subsequent derivatives are known as **higher derivatives**. Second derivatives in particular occur frequently in engineering contexts.



Prerequisites

Before starting this Section you should ...

- be able to differentiate standard functions




Learning Outcomes

On completion you should be able to ...

- obtain higher derivatives

1. The derivative of a derivative

You have already learnt how to calculate the derivative of a function using a table of derivatives and applying some basic rules. By differentiating the function, $y(x)$, we obtain the derivative, $\frac{dy}{dx}$. By repeating the process we can obtain higher derivatives.



Example 7

Calculate the first, second and third derivatives of $y = x^4 + 6x^2$.

Solution

The first derivative is $\frac{dy}{dx}$:

$$\text{first derivative} = 4x^3 + 12x$$

To obtain the second derivative we differentiate the first derivative.

$$\text{second derivative} = 12x^2 + 12$$

The third derivative is found by differentiating the second derivative.

$$\text{third derivative} = 24x + 0 = 24x$$

2. Notation for derivatives

Just as there is a notation for the first derivative so there is a similar notation for higher derivatives. Consider the function, $y(x)$. We know that the first derivative is $\frac{dy}{dx}$ or $\frac{d}{dx}(y)$ which is the instruction to differentiate the function $y(x)$. The second derivative is calculated by differentiating the first derivative, that is

$$\text{second derivative} = \frac{d}{dx} \left(\frac{dy}{dx} \right)$$

So, using a fairly obvious adaptation of our derivative notation, the second derivative is denoted by $\frac{d^2y}{dx^2}$ and is read as 'dee two y by dee x squared'. This is often written more concisely as y'' .

In similar manner, the third derivative is denoted by $\frac{d^3y}{dx^3}$ or y''' and so on. So, referring to Example 6 we could have written

$$\begin{aligned} \text{first derivative} &= \frac{dy}{dx} = 4x^3 + 12x \\ \text{second derivative} &= \frac{d^2y}{dx^2} = 12x^2 + 12 \\ \text{third derivative} &= \frac{d^3y}{dx^3} = 24x \end{aligned}$$



Key Point 7

If $y = y(x)$ then its first, second and third derivatives are denoted by:

$$\frac{dy}{dx} \quad \frac{d^2y}{dx^2} \quad \frac{d^3y}{dx^3}$$

or

$$y' \quad y'' \quad y'''$$

In most examples we use x to denote the independent variable and y the dependent variable. However, in many applications, time t is the independent variable. In this case a special notation is used for derivatives. Derivatives with respect to t are often indicated using a **dot** notation, so $\frac{dy}{dt}$ can be written as \dot{y} , pronounced 'y dot'. Similarly, a second derivative with respect to t can be written as \ddot{y} , pronounced 'y double dot'.



Key Point 8

If $y = y(t)$ then

$$\dot{y} \text{ stands for } \frac{dy}{dt}, \quad \ddot{y} \text{ stands for } \frac{d^2y}{dt^2} \text{ etc}$$



Calculate $\frac{d^2y}{dt^2}$ and $\frac{d^3y}{dt^3}$ given $y = e^{2t} + \cos t$.

First find $\frac{dy}{dt}$:

Your solution

Answer

$$\frac{dy}{dt} = 2e^{2t} - \sin t$$

Now obtain the second derivative:

Your solution

$$\frac{d^2y}{dt^2} =$$

Answer

$$4e^{2t} - \cos t$$

Finally, obtain the third derivative:

Your solution

$$\frac{d^3y}{dt^3} = \frac{d}{dt} \left(\frac{d^2y}{dt^2} \right) =$$

Answer

$$8e^{2t} + \sin t$$

Note that in the last Task we could have used the dot notation and written $\dot{y} = 2e^{2t} - \sin t$, $\ddot{y} = 4e^{2t} - \cos t$ and $\ddot{\dot{y}} = 8e^{2t} + \sin t$

We may need to evaluate higher derivatives at specific points. We use an obvious notation.

The second derivative of $y(x)$, evaluated at say, $x = 2$, is written as $\frac{d^2y}{dx^2}(2)$, or more simply as $y''(2)$.

The third derivative evaluated at $x = -1$ is written as $\frac{d^3y}{dx^3}(-1)$ or $y'''(-1)$.



Given $y(x) = 2 \sin x + 3x^2$ find (a) $y'(1)$ (b) $y''(-1)$ (c) $y'''(0)$

First find $y'(x)$, $y''(x)$ and $y'''(x)$:

Your solution

$$y'(x) = \qquad \qquad \qquad y''(x) = \qquad \qquad \qquad y''' =$$

Answer

$$y'(x) = 2 \cos x + 6x \qquad \qquad y''(x) = -2 \sin x + 6 \qquad \qquad y'''(x) = -2 \cos x$$

Now substitute $x = 1$ in $y'(x)$ to obtain $y'(1)$:

Your solution

$$(a) \quad y'(1) =$$

Answer

$$y'(1) = 2 \cos 1 + 6(1) = 7.0806. \text{ Remember, in } \cos 1 \text{ the '1' is 1 radian.}$$

Now find $y''(-1)$:

Your solution

(b) $y''(-1) =$

Answer

$$y''(-1) = -2 \sin(-1) + 6 = 7.6829$$

Finally, find $y'''(0)$:

Your solution

(c) $y'''(0) =$

Answer

$$y'''(0) = -2 \cos 0 = -2.$$

Exercises

- Find $\frac{d^2y}{dx^2}$ where $y(x)$ is defined by:
(a) $3x^2 - e^{2x}$ (b) $\sin 3x + \cos x$ (c) \sqrt{x} (d) $e^x + e^{-x}$ (e) $1 + x + x^2 + \ln x$
- Find $\frac{d^3y}{dx^3}$ where y is given in Exercise 1.
- Calculate $\ddot{y}(1)$ where $y(t)$ is given by:
(a) $t(t^2 + 1)$ (b) $\sin(-2t)$ (c) $2e^t + e^{2t}$ (d) $\frac{1}{t}$ (e) $\cos \frac{t}{2}$
- Calculate $\ddot{y}(-1)$ for the functions given in Exercise 3.

Answers

- (a) $6 - 4e^{2x}$ (b) $-9 \sin 3x - \cos x$ (c) $-\frac{1}{4}x^{-3/2}$ (d) $e^x + e^{-x}$ (e) $2 - \frac{1}{x^2}$
- (a) $-8e^{2x}$ (b) $-27 \cos 3x + \sin x$ (c) $\frac{3}{8}x^{-5/2}$ (d) $e^x - e^{-x}$ (e) $\frac{2}{x^3}$
- (a) 6 (b) 3.6372 (c) 34.9927 (d) 2 (e) -0.2194
- (a) 6 (b) -3.3292 (c) 1.8184 (d) -6 (e) -0.0599